

On "Pulsar dynamics: magnetic dipole model revisited"

D. P. Barsukov, E. M. Kantor, A. I. Tsygan ^{*},

Ioffe Physical Technical Institute,

Politekhnikeskaya 26, 194021 St. Petersburg, Russia

On November the 17th an article written by G. Lugones and I. Bombaci appeared in astro-ph/0411507. The authors reconsider a pulsar braking due to magneto-dipolar deceleration. The authors made remark that it is necessary to take into account the time dependence of angular velocity Ω while differentiating phase $\phi = \Omega t$ by time, so $\dot{\phi} = \dot{\Omega}t + \Omega$. But their definition of the angular velocity is not conventional, usually the angular velocity is defined as $\dot{\phi}$. Their analysis shows that the polar magnetic field strength B_p inferred from timing observations may be up to a factor 4 larger than currently expected and braking index is also different. This result is not correct. They did not use this expression ($\dot{\phi} = \dot{\Omega}t + \Omega$) while calculating the rotational energy of the pulsar $E = I\dot{\phi}^2/2$ (the formula which was used by the authors is $E = I\Omega^2/2$).

Use of correct formula for the rotational energy leads to the following differential equation for the pulsar deceleration:

$$\dot{\phi}\ddot{\phi} = -K(\dot{\phi}^4 + \ddot{\phi}^2) \quad (1)$$

Neglecting the second term (which is small in comparison to the first one) in the right hand side of this equation, we can rewrite it in traditional form:

$$\ddot{\phi} = -K\dot{\phi}^3 \quad (2)$$

and we arrive at the conventional values for the magnetic fields and braking index equal to 3.

^{*}*e-mail:* tsygan@astro.ioffe.rssi.ru